

# POSTPIL®

Effective and convenient 1-pill solution for emergency contraceptive

EASY-TO-USE · EASY-TO-FIND · EFFECTIVE · SAFE



## WHAT IS POSTPIL®

- **Emergency contraceptive** to use after unprotected intercourse or in case of known or suspected contraceptive failure
- **Package contains 1 tablet of levonorgestrel 1.5 mg**
- **Conforms to Eur. Pharmacopeia** and manufactured at an EU cGMP-approved facility
- **Not an abortion pill**; does not terminate established pregnancy
- **Does not protect against sexually transmitted infections (STIs)**
- **Debunking common myths:**
  - No health risks with repeated uses
  - Does not affect fertility
  - Does not increase the risk of ectopic pregnancy, cancer or cause long term side effects

## WHO CAN USE POSTPIL®

- **Women of all reproductive ages who want to avoid pregnancy**, including adolescents, women who are breastfeeding, or women with history of ectopic pregnancy
- **No absolute contraindications**, except hypersensitivity to levonorgestrel

## WHY USE POSTPIL®

- **WHO-recommended regimen** to prevent pregnancy
- **Meets European standards and certifications** for quality control of medicines and production processes
- **1-pill formulation** is easier to follow correctly than 2-pill formulation
- **High-quality yet affordable product**

### ALSO AVAILABLE FROM WOMANCARE GLOBAL



#### MISOMIFE-FEM® COMBO

Combi pack of 1 mifepristone 200mg tablet and 4 misoprostol 200mcg tablet



#### MISO-MIFE®

Combi pack of 4 misoprostol 200mcg tablets



#### IPAS MVAPlus

Uterine evacuation device

# POSTPIL®

Effective and convenient 1-pill solution for emergency contraceptive

EASY-TO-USE · EASY-TO-FIND · EFFECTIVE · SAFE



## WHEN TO USE POSTPIL®

- **Can be used at any time during the menstrual cycle for the following situations:**
  - Desire to avoid pregnancy
  - Nonconsensual or forced unprotected sex
  - Possible contraceptive failure or incorrect use
  - Concern about the withdrawal method
  - Uncertainty about the 'safe period'
- **Can be used if already administered** in the same menstrual cycle

## HOW POSTPIL® WORKS

- **Disrupts ovulation and thickens the cervical mucus**, which inhibits sperm mobility and prevents fertilization
- **No examinations or laboratory tests needed<sup>1</sup>**
- **Reduces risk of pregnancy by 80%-90%**
- **Side effects may include:** nausea, lower abdominal pain, fatigue, headache, dizziness, breast tenderness, delay of menses (>7 days)

## HOW TO USE POSTPIL®

- **To be used as soon as possible up to 3 days (72 hours)** after unprotected intercourse for best results. Postpil® is more effective the sooner it is taken after unprotected sex.
- **Dosage should not be taken more than once within 24 hours**, except if the woman vomits within 2 hours after use
- **Risk of pregnancy if repeated unprotected intercourse** within 24 hours after taking the Postpil®

ALSO AVAILABLE FROM  
WOMANCARE GLOBAL



### MISOMIFE-FEM® COMBO

Combi pack of 1 mifepristone 200mg tablet and 4 misoprostol 200mcg tablet



### MISO-MIFE®

Combi pack of 4 misoprostol 200mcg tablets



### IPAS MVAPlus

Uterine evacuation device

<sup>1</sup> Rodrigues, Isabel & Grou, Fabienne & Joly, Jacques. (2001). Effectiveness of emergency contraceptive pills between 72 and 120 hours after unprotected sexual intercourse. American journal of obstetrics and gynecology. 184. 531-7. 10.1067/mob.2001.1111102. 1