#### LEVOPLANT LEVOPLANT (h) LEVOPLANT Subdermal contraceptive implant LE VOPLANT . Contraceptive Implant Levoplant<sup>™</sup> is a WHO Prequalifed subdermal nplant(II) LE VOPLANT contraceptive implant effective for up to 3 years. ey are used for a Sino-implant (II It is a safe, highly convenient and acceptable CHARACTERISTICS

High quality<sup>1</sup>, affordable, contraceptive implant that is composed of two thin and flexible silicone rods

WHO pregualified contraceptive implant<sup>2</sup>

Contains 75 mg of levonorgestrel in each 60 month

# EFFICACY

- More than 99% effective in a 3 year period – less than 1 pregnancy per 100 women over duration of use<sup>2</sup>
  - \*For heavier women, the effectiveness of Levoplant<sup>™</sup> may decrease near the end of the duration. Earlier replacement can be discussed.

LE VOPLANT .

rod (150 mg total)

(five year) shelf life

# MECHANISM OF ACTION

- Prevents the release of the egg
- Thickens mucus in the cervical canal to prevent sperm reaching and fertilizing the egg

### ADMINISTRATION

- Inserted just under the skin of a woman's upper arm by a trained healthcare provider using a disposable trocar
- Simple insertion and removal takes a few minutes in the clinical setting with local anaesthetic

## BENEFITS

- Estrogen free: avoids estrogen-related side effects that may be experienced with combined hormonal contraceptive methods
- Action to prevent pregnancy is not required once the implant is inserted.
- The implant can be removed at any time at the woman's request and there is no delay in return to fertility
- Almost all women of any age, including those who have just given birth, are breastfeeding, or women who have had an abortion can use Levoplant™\*

\*if they are medically eligible

### SIDE EFFECTS

Irregular bleeding patterns may occur but are not harmful. Abdominal pain, weight changes, breast tenderness, mood changes, migraine can also be experienced.

## CONTRAINDICATIONS

Unexplained vaginal bleeding, severe liver cirrhosis or liver tumor, history of breast cancer or current deep blood clots.

Steiner MJ, et al. Randomized trial to evaluate contraceptive efficacy, safety and acceptability of a two-rod contraceptive implant over 4 years in the Dominican Republic. 1.

Contracept X. (2019) 1:100006. World Health Organization Department of Sexual and Reproductive Health and Research (WHO/SRH) and Johns Hopkins Bloomberg School of Public Health/ Center for 2 Communication Programs (CCP), Knowledge SUCCESS. Family Planning: A Global Handbook for Providers 4th edition, 2022

WomanCare Global Email: contact@dktwomancare.org www.dktwomancare.org

