

MISO-FEM®

A versatile product with multiple applications

HIGH-QUALITY · COST-EFFECTIVE · EASY-TO-USE



HOW IT WORKS

- **Stimulates the smooth muscle** to cause uterine contractions
- **Softens the cervix** through disintegration and dissolution of collagen
- **Dilates blood vessels** to reduce blood pressure in the uterus
- **Can be administered orally, buccally, or vaginally**

OBSTETRIC AND GYNECOLOGICAL USES

- **Medical abortion:** effective both alone (87%) and in combination with mifepristone (96%)¹ where allowed
- **Miscarriage management:** effective and highly acceptable for treating early pregnancy loss or incomplete abortion
- **Cervical ripening:** effective ripening agent prior to first-trimester aspiration abortions. Also used prior to endometrial biopsy or IUD insertion
- **Labor induction:** highly effective for inducing vaginal delivery within 24 hours
- **Postpartum hemorrhage (PPH):** used for prevention and treatment of PPH

Misoprostol compares favorably to oxytocin for PPH because it is inexpensive, thermostable, has low side-effects rates, and does not require injection²

1. Kapp, N., & Lohr, P. A. (2020). Modern methods to induce abortion: Safety, efficacy and choice. Best Practice & Research Clinical Obstetrics & Gynaecology, 63, 37-44. doi: 10.1016/j.bpobgyn.2019.11.008

2. Allen, R., & O'Brien, B. M. (2009). Uses of misoprostol in obstetrics and gynecology. Reviews in obstetrics & gynecology, 2(3), 159-168.

ALSO AVAILABLE FROM
WOMANCARE GLOBAL



MISOMIFE-FEM® COMBO

For uterine evacuation
Combi pack of 1 mifepristone 200 mg tablet
and 4 misoprostol 200 mcg tablets
Product Code: COMBO



POSTPIL®

Pack of 1 levonorgestrel 1.5 mg tablet
Product Code: EC1



IPAS MVA PLUS® ASPIRATOR

For uterine evacuation
Product Code: PLUS

MISO-FEM®

A versatile product with multiple applications

HIGH QUALITY · COST-EFFECTIVE · EASY-TO-USE




INSTRUCTIONS FOR UTERINE EVACUATION: STEP 1

Administer 4 misoprostol pills:

- **Buccally**, between your cheek and bottom gums for 30 minutes, then swallow with water
- **Sublingually**, under your tongue for 30 minutes, then swallow with water

Suggestions for use:

- Have someone with you for support
- Before taking the pills, take a drink of water to moisten your mouth
- Do not eat or drink anything once you take the pills
- You may experience cramps and bleeding
- Expect to pass clots of blood and tissue



INSTRUCTIONS FOR UTERINE EVACUATION: STEPS 2 & 3

- Wait 3-4 hours
- Repeat step 1 with another 4 pills
- Wait 3-4 hours
- Repeat step 1 with another 4 pills

ALSO AVAILABLE FROM WOMANCARE GLOBAL



IPAS DOUBLE VALVE ASPIRATOR

For uterine evacuation
Product Code: DVS-10 or DVS-SU



IPAS EASYGRIP® CANNULAE

For uterine evacuation
Product Code: SR4 to SR10, and SR12 or SR4-SU to SR10-SU, and SR12-SU



FLEXIBLE KARMAN CANNULAE

For uterine evacuation
Product Code: FC4 to FC10, and FC12