

PRODUCT CODE: COMBO

MISOMIFE-FEM® COMBO

Pack of 1 mifepristone 200 mg tablet and 4 misoprostol 200 mcg tablets

AN EFFECTIVE, HIGH-QUALITY, SAFE, AND DISCREET
OPTION FOR UTERINE EVACUATION

WHAT IS MISOMIFE-FEM® COMBO

- **Mifepristone** 200 mg (1 tablet) stops the action of the hormone that sustains pregnancy (progesterone). This causes the embryo to detach from the uterus and the pregnancy to stop growing
- **Misoprostol** 800 mcg (4 tablets) causes contraction, to push out the contents of the uterus

OBSTETRIC AND GYNECOLOGICAL USES

Ensures complete uterine evacuation:

- For first-trimester abortion or miscarriage
- When therapeutic abortion is required for a pregnancy that is 9 weeks or less
- For treatment of postpartum hemorrhage / Post Abortion Care (PAC)
- To prevent a hysterectomy and shorten the duration of conservative management of placenta accreta



- MORE EFFECTIVE THAN USING MISOPROSTOL ALONE¹
- SAFE METHOD FOR UTERINE EVACUATION
- CAN BE USED OUTSIDE THE HOSPITAL, EVEN AT HOME (WHERE ALLOWED)
- DISCREET AND RESPECTFUL OF THE WOMAN'S PRIVACY
- AVAILABLE FROM PHARMACY (WHERE ALLOWED)
- PREVENTS MATERNAL DEATHS BY AVERTING UNSAFE ABORTION
- REGIMEN RECOMMENDED BY THE WHO

ALSO AVAILABLE FROM
WOMANCARE GLOBAL



MISO-FEM®

For uterine evacuation

Pack of 4 misoprostol 200 mcg tablets


Product Code: MISO



POSTPIL®

Pack of 1 levonorgestrel 1.5 mg tablet

Product Code: EC1



IPAS MVA PLUS® ASPIRATOR

For uterine evacuation

Product Code: PLUS

1. Kapp, N., & Lohr, P. A. (2020). Modern methods to induce abortion: Safety, efficacy and choice. Best Practice & Research Clinical Obstetrics & Gynaecology, 63, 37-44. doi: 10.1016/j.bpobgyn.2019.11.008

MISOMIFE-FEM® COMBO

Pack of 1 mifepristone 200 mg tablet and 4 misoprostol 200 mcg tablets

AN EFFECTIVE, HIGH-QUALITY, SAFE, AND DISCREET
OPTION FOR UTERINE EVACUATION




INSTRUCTIONS FOR UTERINE EVACUATION: STEP 1

Swallow mifepristone tablet with water

- Patient should have someone with them for support
- They should eat something light (e.g. crackers) or take a drug to prevent vomiting (ask for an "anti-emetic" drug)

Wait for 36 hours after taking misoprostol tablet

- Most women do not feel any symptom during this period
- Some may feel like vomiting or have slight bleeding




INSTRUCTIONS FOR UTERINE EVACUATION: STEP 2

Administer 4 tablets of 200 mcg misoprostol (800 mcg total)

- **Buccally**, between your cheek and bottom gums for 30 minutes then swallow with water
- **Sublingually**, under your tongue for 30 minutes then swallow with water

What to expect:

- Clients should expect to have painful cramps and bleeding
- They should expect to pass clots of blood and tissue



ALSO AVAILABLE FROM WOMANCARE GLOBAL



INJECTA-FEM

Intramuscular contraceptive injection
Medroxyprogesterone acetate 150 mg/ml
Product Code: INJECTA



LEVOPLANT™

WHO-prequalified
2-rod hormonal implant
Product Code: LEVO



UTERINE PELVIC MODEL

For uterine evacuation training
Product Code: UPM