


POSTPIL®

Effective and convenient 1-pill solution for emergency contraceptive

EASY-TO-USE · EASY-TO-FIND · EFFECTIVE · SAFE



WHAT IS POSTPIL®

- **Emergency contraceptive** to use after unprotected intercourse or in case of known or suspected contraceptive failure
- **Package contains 1 tablet of levonorgestrel 1.5 mg**
- **Conforms to Eur. Pharmacopeia** and manufactured at an EU cGMP-approved facility
- **Not an abortion pill**; does not terminate established pregnancy
- **Does not protect against sexually transmitted infections (STIs)**
- **Debunking common myths:**
 - No health risks with repeated uses
 - Does not affect fertility
 - Does not increase the risk of ectopic pregnancy, cancer or cause long term side effects

WHO CAN USE POSTPIL®

- **Women of all reproductive ages who want to avoid pregnancy**, including adolescents, women who are breastfeeding, or women with history of ectopic pregnancy
- **No absolute contraindications**, except hypersensitivity to levonorgestrel

WHY USE POSTPIL®

- **WHO-recommended regimen** to prevent pregnancy
- **Meets European standards and certifications** for quality control of medicines and production processes
- **1-pill formulation** is easier to follow correctly than 2-pill formulation
- **High-quality yet affordable product**

ALSO AVAILABLE FROM WOMANCARE GLOBAL



MISOMIFE-FEM® COMBO

Combi pack of 1 mifepristone 200mg tablet and 4 misoprostol 200mcg tablet



MISO-MIFE®

Combi pack of 4 misoprostol 200mcg tablets




IPAS MVAPlus

Uterine evacuation device

POSTPIL®

Effective and convenient 1-pill solution for emergency contraceptive

EASY-TO-USE · EASY-TO-FIND · EFFECTIVE · SAFE



WHEN TO USE POSTPIL®

- **Can be used at any time during the menstrual cycle for the following situations:**
 - Desire to avoid pregnancy
 - Nonconsensual or forced unprotected sex
 - Possible contraceptive failure or incorrect use
 - Concern about the withdrawal method
 - Uncertainty about the 'safe period'
- **Can be used if already administered** in the same menstrual cycle

HOW POSTPIL® WORKS

- **Disrupts ovulation and thickens the cervical mucus**, which inhibits sperm mobility and prevents fertilization
- **No examinations or laboratory tests needed**¹
- **Reduces risk of pregnancy by 80%-90%**
- **Side effects may include:** nausea, lower abdominal pain, fatigue, headache, dizziness, breast tenderness, delay of menses (>7 days)

HOW TO USE POSTPIL®

- **To be used as soon as possible up to 5 days (120 hours)** after unprotected intercourse for best results. Postpil® is more effective the sooner it is taken after unprotected sex.
- **Dosage should not be taken more than once within 24 hours**, except if the woman vomits within 2 hours after use
- **Risk of pregnancy if repeated unprotected intercourse** within 24 hours after taking the Postpil®

ALSO AVAILABLE FROM
WOMANCARE GLOBAL



MISOMIFE-FEM® COMBO

Combi pack of 1 mifepristone 200mg tablet and 4 misoprostol 200mcg tablet



MISO-MIFE®

Combi pack of 4 misoprostol 200mcg tablets



IPAS MVAPlus

Uterine evacuation device

¹ Rodrigues, Isabel & Grou, Fabienne & Joly, Jacques. (2001). Effectiveness of emergency contraceptive pills between 72 and 120 hours after unprotected sexual intercourse. American journal of obstetrics and gynecology. 184. 531-7. 10.1067/mob.2001.111102. 1